

University of Dundee

Duration of dual antiplatelet therapy in acute coronary syndrome

Wilson, Simon John; Newby, David E.; Dawson, Dana; Irving, John; Berry, Colin

Published in:
Heart

DOI:
[10.1136/heartjnl-2016-309871](https://doi.org/10.1136/heartjnl-2016-309871)

Publication date:
2017

Licence:
CC BY

Document Version
Publisher's PDF, also known as Version of record

[Link to publication in Discovery Research Portal](#)

Citation for published version (APA):

Wilson, S. J., Newby, D. E., Dawson, D., Irving, J., & Berry, C. (2017). Duration of dual antiplatelet therapy in acute coronary syndrome. *Heart*, 103(8), 573-580. <https://doi.org/10.1136/heartjnl-2016-309871>

General rights

Copyright and moral rights for the publications made accessible in Discovery Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from Discovery Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain.
- You may freely distribute the URL identifying the publication in the public portal.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



OPEN ACCESS

Duration of dual antiplatelet therapy in acute coronary syndrome

Simon John Wilson,¹ David E Newby,¹ Dana Dawson,⁶ John Irving,² Colin Berry³

¹British Heart Foundation Centre for Cardiovascular Science, New Royal Infirmary of Edinburgh, Edinburgh, UK

²Department of Cardiology, Ninewells Hospital, Dundee, UK

³British Heart Foundation Glasgow Cardiovascular Research Centre, University of Glasgow, Glasgow, UK

⁶Department of Cardiovascular Medicine, School of Medicine, Medical Sciences and Nutrition, University of Aberdeen, Aberdeen

Correspondence to

Professor David E Newby, BHF Centre for Cardiovascular Science, Room SU314, Chancellor's Building, University of Edinburgh, 49 Little France Crescent, Edinburgh, EH16 4SA, UK; d.e.newby@ed.ac.uk

Received 28 October 2016

Revised 02 January 2017

Accepted 03 January 2017

ABSTRACT

Despite a large volume of evidence supporting the use of dual antiplatelet therapy in patients with acute coronary syndrome, there remains major uncertainty regarding the optimal duration of therapy. Clinical trials have varied markedly in the duration of therapy, both across and within trials. Recent systematic reviews and meta-analyses suggest that shorter durations of dual antiplatelet therapy are superior because the avoidance of atherothrombotic events is counterbalanced by the greater risks of excess major bleeding with apparent increases in all-cause mortality with longer durations. These findings did not show significant heterogeneity according to whether patients had stable or unstable coronary heart disease. Moreover, the potential hazards and benefits may differ when applied to the general broad population of patients encountered in everyday clinical practice who have markedly higher bleeding and atherothrombotic event rates. Clinicians lack definitive information regarding the duration of therapy in patients with acute coronary syndrome and risk scores do not appear to be sufficiently robust to address these concerns. We believe that there is a pressing need to undertake a broad inclusive safety trial of shorter durations of therapy in real world populations of patients with acute coronary syndrome. The clinical evidence would further inform future research into strategies for personalised medicine.

INTRODUCTION

A ruptured or eroded coronary atherosclerotic plaque is the principal underlying cause of an acute coronary syndrome. The greatest 'at risk' period is during this early phase of plaque instability and healing, with recurrent event rates peaking in the first month. By 3 months, the plaque has usually stabilised, healed and subsequent event rates return to the background rates seen in patients with stable coronary heart disease.¹⁻³ Indeed, beyond 3 months, recurrent events commonly occur on plaques at other sites within the coronary circulation.³ From first principles, the first 3 months is the most critical time for interventions to reduce recurrent cardiovascular events after an acute coronary syndrome (ACS). This is consistent with event rates seen in all clinical trials of patients with acute coronary syndrome: an initial time-varying high event rate that reverts to a consistent linear lower event rate from 3 months onwards (table 1).^{1,2,4,5}

Antiplatelet therapy

In an acute coronary syndrome, thrombus formation occurs under conditions of high shear stress and is principally driven by platelet aggregation (figure 1). This dominance of platelet aggregation

during intracoronary thrombus formation reflects the dramatic effects that antiplatelet therapies have on clinical outcomes (table 2). Aspirin was the first antiplatelet therapy which induced a halving in event rates in patients with acute coronary syndrome:^{6,7} such a large effect size has rarely been surpassed in other domains of cardiology.

Given aspirin's remarkable success, it is perhaps unsurprising that adjunctive antiplatelet therapies have been investigated to build on these benefits, especially as there are multiple mechanisms of platelet activation beyond the cyclo-oxygenase pathway (figure 2). However, as platelets are essential to primary haemostasis, there is a balance between reducing the incidence of future cardiovascular events and causing harm from an increased risk of bleeding. The P2Y₁₂ receptor antagonists are a class of drugs that have gained widespread acceptance since they appear to provide additional thrombotic protection at the expense of modest increases in bleeding. Their use is principally associated with reductions in recurrent myocardial infarction^{1,4,5,8} and in a few trials, reductions in cardiovascular events and mortality.^{5,8} Other antiplatelet therapies (figure 2) are available but have variable net clinical benefit and for the purposes of this review, we will consider only dual antiplatelet therapy (DAPT) with aspirin and P2Y₁₂ receptor antagonism.

Dual antiplatelet therapy

The benefit of dual antiplatelet therapy following an acute coronary syndrome was established by the CURE,¹ COMMIT/CCS-2⁸ and CLARITY-TIMI 28⁹ trials. Combined aspirin and clopidogrel therapy reduced the 1-year incidence of cardiovascular events by approximately 20% compared with aspirin alone. More potent and consistent P2Y₁₂ receptor inhibition with either prasugrel or ticagrelor was superior to clopidogrel in the subsequent TRITON⁴ and PLATO⁵ trials.

The evidence for dual antiplatelet therapy in patients with stable coronary heart disease is less distinct. In the CHARISMA trial,² the addition of clopidogrel to aspirin in patients with established cardiovascular disease or at high risk of clinical atherosclerotic disease did not reduce cardiovascular events and was associated with an increase in severe bleeding. There was, however, a suggestion of improved outcomes in patients with established atherothrombotic disease, particularly those with a history of myocardial infarction. The PEGASUS-TIMI 54¹⁰ trial compared aspirin monotherapy to a combination of aspirin and ticagrelor in patients with a previous myocardial infarction and at least one additional high-risk factor. At a



CrossMark

To cite: Wilson SJ, Newby DE, Dawson D, *et al.* Heart Published Online First: [please include Day Month Year]. doi:10.1136/heartjnl-2016-309871

Table 1 Temporal relationship with the clinical benefits of clopidogrel therapy

Time interval (months)	Primary endpoint*		ARR (%)	RRR [95% CIs] (%)	NNT (per month)
	Clopidogrel (%)	Placebo (%)			
CURE trial					
0–1	4.3	5.5	1.2	22 [9, 32]	84
1–3	1.8	2.5	0.8	32 [13, 46]	240
3–6	1.8	1.8	0.0	4 [–27, 27]	5174
6–9	1.3	1.4	0.1	6 [–34, 34]	3171
9–12	1.1	1.3	0.2	14 [–32, 44]	1600
0–12	10.3	12.6	2.4	19	507
CHARISMA trial					
0–28	6.8	7.3	0.5	7*	5591
Subgroup of patients with clinically evident atherosclerotic disease					
0–28†	6.9	7.9	1.0	12	2800

*Primary endpoint—cardiovascular death, myocardial infarction and stroke.

ARR, absolute risk reduction; CI, confidence intervals; CVD, cardiovascular death; MI, myocardial infarction; NNT, number needed to treat; RRR, relative risk reduction.

mean of 33 months, ticagrelor (60 mg) reduced the incidence of cardiovascular death, myocardial infarction or stroke (7.77% vs 9.04%) at the expense of increased thrombolysis in myocardial infarction (TIMI) major bleeding (2.30% vs 1.06%) and a neutral effect on overall mortality. On the basis of these trials, combination antiplatelet therapy would appear to confer only a small ischaemic benefit at the cost of a significant bleeding risk. European¹¹ and North American¹² guidelines therefore do not recommend dual antiplatelet therapy in patients with stable atherothrombotic disease but acknowledge that with careful consideration, combined antiplatelet therapy may be beneficial in some high-risk patients.

Duration of dual antiplatelet therapy: clinical trials

Current European¹³ and North American¹² guidelines advise continuing dual antiplatelet therapy for 1 year following an acute coronary syndrome. These recommendations are made on the basis of early studies^{4 5 14 15} demonstrating a sustained increased

risk of thrombotic complications, including stent thrombosis and spontaneous cardiovascular events, beyond 6 months. However, the greatest absolute reductions in cardiovascular events with dual antiplatelet therapy are seen in the first 3 months (table 1) and since these studies, advances in drug-eluting stent technology have led to a substantially reduced incidence of late (>30 days) and very late (>1 year) stent thrombosis.¹⁶

In recent trials of patients treated with newer generation drug-eluting stents, shorter durations of dual antiplatelet therapy (3 months to 6 months) were non-inferior to 12^{17–22} months or 24²³ months of treatment with regard to either a composite of cardiovascular events or cardiovascular events plus major bleeding. Moreover, all of these trials included patients with an acute coronary syndrome (range 23% to 74% of study population) and in those who undertook prespecified subgroup analyses, no heterogeneity in treatment effect between stable and unstable coronary artery disease was observed (figure 3).

Beyond 12 months, there remains a residual risk of local and systemic atherothrombotic complications²⁴ and a number of studies have examined whether extended dual antiplatelet therapy (>12 months) following percutaneous coronary intervention may be beneficial. In the DES-LATE²⁵ and ARCTIC-INTERRUPTION trials,²⁶ prolonged treatment with dual antiplatelet therapy (18–30 months vs 12 months) neither reduced the incidence of cardiovascular events nor increased the risk of major bleeding. Among those patients presenting with an acute coronary syndrome, primary and secondary ischaemic end points did not differ from the global treatment population. In the DAPT trial,²⁷ the largest and only double-blinded study, extended dual antiplatelet therapy (30 months vs 12 months) reduced the risk of major adverse cardiovascular and cerebrovascular events (4.3% vs 5.9%), myocardial infarction (2.1% vs 4.1%) and stent thrombosis (0.4% vs 1.4%) but at a cost of increased moderate or severe bleeding (2.5% vs 1.6%) and a borderline rise in all-cause mortality (2.0% vs 1.0%; $p=0.05$). Treatment effect did not differ between patients with or without a history of myocardial infarction for any of the co-primary end points including bleeding (figure 3).




Figure 1 Immunofluorescent staining of human thrombus (platelets, green; fibrin, red) formed at high shear stress.

Table 2 Major trials of antiplatelet agents in acute coronary syndrome \pm unstable angina

Antiplatelet agent	Mechanism of action	Trial	Comparison	(Primary) end point	Risk reduction (time point)
Aspirin	COX-1 inhibitor	The ISIS-2 collaborators (ISIS-2), 1979 The Risk Group, 1990 ⁵¹ Lewis <i>et al</i> , 1983 ⁵² Cairns <i>et al</i> , 1985 ⁵³	aspirin versus placebo aspirin versus placebo aspirin versus placebo aspirin versus placebo	Vascular mortality MI or death MI or death MI or death	23% (5 weeks) 74% (3 months) 51% (3 months) 51% (2 years)
Ticlopidine	P2Y12 antagonist	Scrutinio <i>et al</i> (STAMI), 2001	aspirin versus ticlopidine	Death, MI, stroke or angina	ns (6 months)
Clopidogrel	P2Y12 antagonist	Bertrand <i>et al</i> (CLASSICS), 2000 Yusuf <i>et al</i> (CURE), 2001 The COMMIT Group (COMMIT), 2005 Sabatine <i>et al</i> (CLARITY), 2005	aspirin + clopidogrel versus aspirin + ticlopidine aspirin + clopidogrel versus aspirin aspirin + clopidogrel versus aspirin aspirin + clopidogrel versus aspirin	Cardiac death, MI, or TLR CV death, MI or stroke Death, MI or stroke CV death, MI or urgent TVR	ns (30 days) 20% (1 year) 9% (discharge or 28 days) 20% (30 days)
Prasugrel	P2Y12 antagonist	Wiviott <i>et al</i> (TRITON), 2007	aspirin + prasugrel versus aspirin + clopidogrel	CV death, MI or stroke	19% (1 year)
Ticagrelor	P2Y12 antagonist	Steg <i>et al</i> (PLATO), 2010	aspirin + ticagrelor versus aspirin + clopidogrel	CV death, MI or stroke	13% (1 year)
Dipyridamole	PDE inhibitor	The PARIS Research Group (PARIS-1), 1980 White <i>et al</i> , 1995 ⁵⁴	aspirin + dipyridamole versus aspirin aspirin + dipyridamole versus aspirin	Cardiac death or MI Prevention of late reocclusion	ns (20 months) ns (1 year)
Cilostazol	PDE inhibitor	Lee <i>et al</i> (DECLARE-LONG II), 2011	Cilostazol + standard care versus standard care	In-stent late loss	18% (8 months)
Abciximab	GPIIb/IIIa inhibitor	The EPIC Investigators (EPIC), 1994 The CAPTURE Investigators (CAPTURE), 1997 Kastrati <i>et al</i> (ISAR-REACT 2), 2006 Simoons <i>et al</i> (GUSTO IV-ACS), 2001 Kastrati <i>et al</i> (IASR-REACT 4), 2011	12-hour infusion versus placebo 24-hour infusion versus placebo 12-hour infusion versus placebo 24-hour or 48-hour infusion versus placebo Abciximab + heparin versus bivalirudin	Death, MI or urgent revascularisation Death, MI or urgent revascularisation Death, MI or urgent TVR Death, MI or urgent TVR Death or MI Death, MI, urgent TVR or major bleeding	35% (30 days) 29% (30 days) 24% (30 days) ns (30 days) ns (30 days)
Eptifibatide	GPIIb/IIIa inhibitor	The PURSUIT Investigators (PURSUIT), 1998	Bolus and 72-hour infusion versus placebo	Death or MI	10% (30 days)
Tirofiban	GPIIb/IIIa inhibitor	The PRISM investigators (PRISM), 1998 The PRISM investigators (PRISM-PLUS), 1998	Bolus and 48-hour infusion versus placebo Bolus and 72-hour infusion versus placebo	Death, MI, refractory ischaemia or readmission for UA Death, MI or refractory ischaemia	ns (30 days) 17% (30 days)
Vorapaxar	PAR-1 antagonist	Tricoci <i>et al</i> (TRACER), 2012	vorapaxar + standard care versus standard care	CV, death, MI, readmission with ischaemia, urgent revascularisation	ns (median 502 days)

COX-1, cyclo-oxygenase-1; CV, cardiovascular; GP, glycoprotein; MI, myocardial infarction; PAR-1, protease-activated receptor-1; PDE, phosphodiesterase; TLR, target lesion revascularisation; TVR, target vessel revascularisation; UA, unstable angina.

Duration of dual antiplatelet therapy: systematic reviews and meta-analyses

Meta-analyses of trials using dual antiplatelet therapy in patients receiving intracoronary stents have compared short (3–6 months), 12-month and prolonged (>12 months) durations of therapy.^{28–29} Longer treatment periods reduced the incidence of myocardial infarction and stent thrombosis but at a cost of increased major bleeding and with a tendency to increase overall mortality because of an increase in non-cardiovascular death. However, the majority of patients included in these analyses had stable coronary artery disease and few patients with acute coronary syndrome were treated with ≤ 6 months of dual antiplatelet therapy. In a recent meta-analysis³⁰ that included only patients with a history of acute coronary syndrome, prolonged dual antiplatelet therapy reduced the risk of cardiovascular death (RR 0.85; 95% CI: 0.74 to 0.98, $p=0.03$) without an increase in non-cardiovascular death (RR 1.03, 95% CI 0.86 to 1.23; $p=0.76$) or all-cause mortality (RR 0.92, 95% CI 0.83 to 1.03).

Atherothrombotic risk

The optimal duration of dual antiplatelet therapy is dependent on the balance between preventing future atherothrombotic events

and the increased risk of bleeding from continued treatment. Following an acute coronary syndrome, predictors of atherothrombotic risk include ST deviation, diabetes mellitus, smoking status, left ventricular ejection fraction, stent type, number of stents and complexity of coronary artery disease. Subgroup analyses have attempted to identify if any of these factors influence outcomes with regard to duration of dual antiplatelet treatment.

In the EXCELLENT trial,²² there was a threefold increase in the incidence of cardiovascular events ($p<0.001$ for interaction) among patients with diabetes mellitus treated with 6 months as compared with 12 months of dual antiplatelet therapy. In the ISAR-SAFE trial,²⁰ rates of death, myocardial infarction, stent thrombosis, stroke or TIMI major bleeding tended to be higher in patients aged <67 years and lower in patients aged ≥ 67 years with 6 months compared with 12 months of dual antiplatelet treatment ($p=0.03$). These differences were driven by ischaemic complications rather than bleeding events. In the DAPT trial, men were nearly five times less likely to suffer from stent thrombosis if dual antiplatelet therapy was extended beyond 12 months ($p=0.04$) and the reduction in major adverse cardiovascular and cerebrovascular events with prolonged therapy (30 months) was greater in those treated with prasugrel (vs clopidogrel; $p=0.03$)




Figure 2 Platelet activation pathways and sites targeted by current and novel antiplatelet agents. Arachidonic acid; ADP, adenosine diphosphate; c, cyclic; Ca²⁺, calcium; AMP, adenosinemonophosphate; COX-1, cyclo-oxygenase-1; DAG, diacylglycerol; GMP, guanosine monophosphate; GP, glycoprotein; IP3: inositol trisphosphate; PAR, protease activated receptor; PDE, phosphodiesterase; PI3K, phosphatidylinositol 3-kinase; PIP2: phosphatidylinositol 4,5-bisphosphate; PLC, phospholipase C; TP, thromboxane receptor; TRPC, transient receptor potential channel; TXA₂, thromboxane A₂; vWF, von Willebrand factor.

or a first-generation drug-eluting stent ($p=0.048$). Similar trends were observed in DES-LATE²⁵ and ARCTIC-INTERRUPTION.²⁶

Audit of the British Cardiovascular Society Intervention database indicates that in 2013/2014, a half of all percutaneous coronary intervention procedures were associated with residual disease (≥ 1 stenosis of $>50\%$ severity), fulfilling the criteria for incomplete revascularisation. Patients with acute coronary syndrome and incomplete revascularisation have a residual burden of coronary disease that is a substrate for recurrent plaque rupture, coronary thrombosis and future cardiac events.^{31–32} Prolonged dual antiplatelet treatment may mitigate this risk but whether this translates to a more favourable risk-to-benefit balance for patients with incomplete revascularisation remains an area for future research.

Bleeding and total mortality

Large registries and trials have shown that major bleeding is associated with an increase in mortality that could potentially negate the benefits of dual antiplatelet therapy in acute coronary syndrome.^{33–36} Importantly, these bleeding risks are not confined to the initial hospitalisation phase.^{33–35} The association between bleeding and mortality has been a consistent feature of acute coronary syndrome trials irrespective of whether the intervention being assessed and improvements in outcome are seen with interventions that are associated with a lower bleeding risk. For example, in the OASIS-5 trial,³³ fondaparinux had

similar antithrombotic benefits to enoxaparin but was associated with lower rates of major bleeding and marked reductions in all-cause mortality. Similar benefits have also been reported for randomised controlled trials of arterial access sites in patients treated with an invasive strategy for either ST-segment³⁷ or non-ST-segment³⁸ myocardial infarction. Again, because radial artery access was associated with less bleeding, overall all-cause mortality was lower.^{37–38} There have been various mechanisms proposed for the link between bleeding and mortality that include rebound hypercoagulability, discontinuation of anti-thrombotic treatments, inflammation and ischaemia.³⁹ The European Society of Cardiology Working Group on Thrombosis has called for clinical trials to address bleeding in acute coronary syndrome including the exploration of the duration of dual antiplatelet therapy.³⁹

Duration uncertainty

Currently there are variations in local and regional dual antiplatelet therapy practices that are confusing for patients, primary care physicians and cardiologists. Indeed, while European¹³ and North American¹² guidelines recommend dual antiplatelet therapy for 12 months after an acute coronary syndrome, both acknowledge that shorter or longer durations may be appropriate. Duration of therapy is seen as a major priority for future research by numerous national and international guideline committees as well as having considerable financial implications, especially for the latest generation of P2Y12 receptor antagonists.

Hazard ratios for the composite primary end-point in patents presenting with and without ACS




Figure 3 Hazard ratios for the composite primary end-point from sub-group analyses of patents presenting with and without an acute coronary syndrome. EXCELLENT trial (n=1443), 6 vs 12 months, patients presenting with ACS = 52% of the study population; PRODIGY trial (n=2013), 6 vs 24 months, patients presenting with ACS subgroup = 74% of study population; ISAR-SAFE trial (n=4000), 6 vs 12 months, patients presenting with ACS = 40% of the study population; OPTIMIZE trial (n=3119), 3 vs 12 months, patients presenting with ACS = 37% of the study population; ARCTIC INTERRUPTION (n=1259), 12 vs 30 months, patients presenting with ACS = 26% of the study population; DES-LATE (n=5045), 12 vs 24 months, patients presenting with ACS = 61% of the study population and DAPT (n=9961), 12 vs 30 months, patients presenting with ACS = 43% of the study population.

However, major pharmaceutical companies have to date not funded trials comparing shorter (<12 months) durations of dual antiplatelet therapy, since, arguably, it may not be in their commercial interest to do so.

For clinicians and healthcare providers, there remains much uncertainty regarding the default duration of dual antiplatelet therapy for most patients with acute coronary syndrome. Current guidelines are largely based on evidence that predates potentially important technological advances, including second-generation drug-eluting stents, while in recent trials, only a minority of patients presented with an acute coronary syndrome and many of these studies were underpowered to detect differences due to low event rates. Selected populations included in randomised controlled trials have lower rates of bleeding and non-cardiovascular death than the general population (table 3), since patients with any history of bleeding or major comorbidity were specifically excluded from such trials. On the other hand, shorter durations of dual antiplatelet therapy may expose medically managed patients (such as due to complex disease) to an increased risk of atherothrombotic events. This is because stenting in acute coronary syndrome confers protection against atherothrombotic

events^{40 41} and in the vast majority of clinical trials, only patients who underwent stent implantation were included. There is therefore a major concern that the evidence to date has been extrapolated to a broader population with a higher risk of both atherothrombotic events and adverse outcomes from bleeding. Accordingly, this has left uncertainty as to the relative benefits and risks of one period of treatment versus another in real-world patients with acute coronary syndrome.

Increased thrombin generation⁴² and platelet reactivity⁴³ have been demonstrated for up to 2 years following a plaque rupture myocardial infarction. This may explain potentially more favourable outcomes with longer durations of dual antiplatelet therapy in patients with acute coronary syndrome as compared to those with stable ischaemic heart disease undergoing percutaneous coronary intervention. However, switching from dual antiplatelet treatment to monotherapy (usually aspirin alone) is associated with a rebound prothrombotic effect, especially with regard to an excess of stent thrombosis.^{10 27} In the DAPT trial,²⁷ this phenomenon occurred irrespective of the timing of switching to monotherapy. Thus, unless dual antiplatelet therapy is continued indefinitely, there will remain a small

Table 3 Temporal relationship with the clinical benefits of clopidogrel therapy

Outcome	CURE (clopidogrel)	TRITON (prasugrel)	TRILOGY (prasugrel)	PLATO (ticagrelor)	Randomised controlled trial average	Scotland 2006–2010* (clopidogrel)
All-cause mortality	5.8%	3.0%	8.3%	4.5%	5.40%	25.4%
Cardiovascular death	5.1%	2.1%	6.6%	4.0%	4.45%	17.8%
Non-cardiovascular death	0.7%	0.9%	1.7%	0.5%	0.95%	7.6%
Fatal bleeding	0.2%	0.4%	0.2%	0.3%	0.28%	0.9%

*From Information and Statistics Division of NHS Scotland.

Patient Characteristics

Age

Diabetes Mellitus ☒

Cigarette Smoking Within Last Two Years ☐

Prior Myocardial Infarction or Percutaneous Coronary Intervention ☒

History of Congestive Heart Failure or Left Ventricular Ejection Fraction < 30% ☐

Index Procedure Characteristics


Myocardial Infarction at Presentation ☒

Stenting of Vein of Graft ☐

Stent Diameter < 3mm ☐

DAPT Score = 2: HIGH
Calculated DAPT Score Shown by Red Arrow

Distribution of DAPT Scores in the DAPT Study



Risk Difference in Events Rates for Continued Thienopyridine Therapy vs. Placebo




Figure 4 DAPT Score calculator for predicting risk/benefit of extending dual antiplatelet therapy from 12 months to 30 months. Available at www.daptstudy.org.

persistent short-term 3-month risk of rebound stent thrombosis and myocardial infarction following the transition from dual to single antiplatelet therapy.

Newer generation P2Y₁₂ inhibitors provide more effective antithrombotic protection than clopidogrel but at the cost of increased bleeding. Given the greater expense of these agents and the marked temporal decline in thrombotic risk that is evident over the first few months, an early 'switch' from ticagrelor or prasugrel to clopidogrel after 1 month to 6 months has been advocated. While evidence from small cohort⁴⁴ and pharmacodynamic studies⁴⁵ suggest such an approach may be safe and reduce bleeding events, an overall lack of data limits any meaningful recommendations. TROPICAL-ACS (NCT 01959451) is an ongoing clinical trial investigating whether a switch to clopidogrel treatment after 1 week of prasugrel is non-inferior to 12 months of standard treatment with prasugrel. Results from this and other similar randomised studies (SWAP-4, NCT 02287909) should provide insights into defining the best strategy for switching between P2Y₁₂ antagonists.

Personalised medicine

Generic recommendations for length of dual antiplatelet therapy derived from a protocolised intervention in clinical trials will inevitably expose some patients to an excessive duration of treatment and disadvantage other patients by withdrawing therapy that protects them from myocardial infarction. The critical clinical question is therefore: can individuals who are more or less likely to benefit from shorter or longer durations of treatment be

identified? While several risk tools have been developed to help determine the future incidence of coronary thrombotic events and major bleeding episodes for an individual, the DAPT Score⁴⁶ and PARIS registry risk Score⁴⁷ were specifically designed to predict medium-term to long-term risks that are directly modified by continuing or interrupting dual antiplatelet therapy.

Using data from the DAPT trial, the DAPT Score was developed to determine the net clinical benefit of extending dual antiplatelet therapy from 12 months to 30 months (figure 4). Thirty-seven candidate variables were considered with 8 included in the final model. In the validation cohort the c-statistic was 0.64 for ischaemia and 0.64 for bleeding. The PARIS registry risk score was similarly developed but was based on prospective observational data. Model discrimination for ischaemia (c-statistic 0.65) and bleeding (c-statistic 0.64) events were comparable to the DAPT score. Thus while these and other similar risk / benefit tools may provide a step forward, they do not as yet offer sufficiently robust predictive value for everyday clinical use.

Variables included in risk tools tend to predict both bleeding and myocardial infarction. High platelet reactivity was originally reported to identify patients at an increased risk of future atherothrombotic events.⁴⁸ However, subsequent studies have failed to show an association between platelet reactivity and atherothrombotic outcomes⁴⁹ or demonstrate that dose adjustment on the basis of on-treatment platelet reactivity is beneficial.⁵⁰ Routine platelet testing is therefore not recommended in patients prescribed P2Y₁₂ inhibitors and is unlikely to deliver a clinically useful prediction score. Further research is needed to define

prediction tools with high discriminatory value in real-world patients, and the potential for net-clinical benefit according to the length of dual anti-platelet therapy.

CONCLUSIONS

It has been 15 years since the CURE trial demonstrated the benefit of dual antiplatelet therapy following an acute coronary syndrome and yet the optimal duration remains uncertain. With regard to thrombotic complications, recent clinical trials and meta-analyses suggest that with newer generation drug-eluting stents, 3 months to 6 months of dual antiplatelet therapy is non-inferior to 12 months of treatment. Prolonged treatment (>12 months) reduces the risk of stent thrombosis, myocardial infarction and possibly cardiovascular death but at the cost of increased major bleeding and with no net mortality benefit. However, these potential hazards and benefits of intervention may differ when applied to the general broad population of patients encountered in everyday clinical practice who have higher bleeding and atherothrombotic event rates.

While ongoing randomised clinical trials may address some of the residual uncertainties in select subgroups, we believe there is a pressing need to undertake a broad inclusive trial of shorter durations of therapy in broad populations of patients with acute coronary syndrome. Such a trial will need to be able to explore specific subgroups, such as those who are medically managed, undergo percutaneous coronary intervention or have coronary artery bypass graft surgery, as well as enable better identification of atherothrombotic and bleeding risks from real world data to inform a more personalised approach to decisions regarding treatment duration.

Acknowledgements DEN is supported by the British Heart Foundation (CH/09/002, RM/13/2/30158, RE/13/3/30183) and is the recipient of a Wellcome Trust Senior Investigator Award (WT103782AIA).

Open Access This is an Open Access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) license, which permits others to distribute, remix, adapt and build upon this work, for commercial use, provided the original work is properly cited. See: <http://creativecommons.org/licenses/by/4.0>

© Article author(s) (or their employer(s) unless otherwise stated in the text of the article) 2017. All rights reserved. No commercial use is permitted unless otherwise expressly granted.

REFERENCES

- Yusuf S, Zhao F, Mehta SR, *et al*; Clopidogrel in Unstable Angina to Prevent Recurrent Events Trial Investigators. Effects of clopidogrel in addition to aspirin in patients with acute coronary syndromes without ST-segment elevation. *N Engl J Med* 2001;345:494–502.
- Bhatt DL, Fox KA, Hacke W, *et al*; CHARISMA Investigators. Clopidogrel and aspirin versus aspirin alone for the prevention of atherothrombotic events. *N Engl J Med* 2006;354:1706–17.
- Stone GW, Maehara A, Lansky AJ, *et al*; PROSPECT Investigators. A prospective natural-history study of coronary atherosclerosis. *N Engl J Med* 2011;364:226–35.
- Wiviott SD, Braunwald E, McCabe CH, *et al*; TRITON-TIMI 38 Investigators. Prasugrel versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2007;357:2001–15.
- Wallentin L, Becker RC, Budaj A, *et al*; PLATO Investigators. Ticagrelor versus clopidogrel in patients with acute coronary syndromes. *N Engl J Med* 2009;361:1045–57.
- Cairns JA, Gent M, Singer J, *et al*. Aspirin, sulfapyrazone, or both in unstable angina. Results of a Canadian multicenter trial. *N Engl J Med* 1985;313:1369–75.
- Lewis HD, Davis JW, Archibald DG, *et al*. Protective effects of aspirin against acute myocardial infarction and death in men with unstable angina. results of a veterans administration cooperative study. *N Engl J Med* 1983;309:396–403.
- Chen ZM, Jiang LX, Chen YP, *et al*; COMMIT (Clopidogrel and Metoprolol in Myocardial Infarction Trial) collaborative group. Addition of clopidogrel to aspirin in 45,852 patients with acute myocardial infarction: randomised placebo-controlled trial. *Lancet* 2005;366:1607–21.
- Sabatine MS, Cannon CP, Gibson CM, *et al*; CLARITY-TIMI 28 Investigators. Addition of clopidogrel to aspirin and fibrinolytic therapy for myocardial infarction with ST-segment elevation. *N Engl J Med* 2005;352:1179–89.
- Bonaca MP, Bhatt DL, Cohen M, *et al*; PEGASUS-TIMI 54 Steering Committee and Investigators. Long-term use of ticagrelor in patients with prior myocardial infarction. *N Engl J Med* 2015;372:1791–800.
- Members TF, Montalescot G, Sechtem U, *et al*. ESC guidelines on the management of stable coronary artery disease. *Eur Heart J* 2013;2013:2949–3003.
- Levine GN, Bates ER, Bittl JA, *et al*. ACC/AHA guideline focused update on duration of dual antiplatelet therapy in patients with coronary artery disease: a report of the American college of cardiology/American heart association task force on clinical practice guidelines: an update of the 2011 ACCF/AHA/SCAI guideline for percutaneous coronary intervention, 2011 ACCF/AHA guideline for coronary artery bypass graft surgery, 2012 ACC/AHA/ACP/AATS/PCNA/SCAI/STS guideline for the diagnosis and management of patients with stable ischemic heart disease, 2013 ACCF/AHA guideline for the management of ST-Elevation myocardial infarction, 2014 AHA/ACC guideline for the management of patients with Non-ST-Elevation Acute Coronary Syndromes, and 2014 ACC/AHA Guideline on Perioperative Cardiovascular Evaluation and Management of Patients Undergoing Noncardiac Surgery. *Circulation* 2016;2016:e123–55.
- Roffi M, Patrono C, Collet J-P, *et al*. ESC guidelines for the management of acute coronary syndromes in patients presenting without persistent ST-segment elevation. *Eur Heart J* 2015;2016:267–315.
- Pfisterer M, Brunner-La Rocca HP, Buser PT, *et al*; BASKET-LATE Investigators. Late clinical events after clopidogrel discontinuation may limit the benefit of drug-eluting stents: an observational study of drug-eluting versus bare-metal stents. *J Am Coll Cardiol* 2006;48:2584–91.
- Eisenstein EL, Anstrom KJ, Kong DF, *et al*. Clopidogrel use and long-term clinical outcomes after drug-eluting stent implantation. *JAMA* 2007;297:159–68.
- Palmerini T, Biondi-Zoccai G, Della Riva D, *et al*. Clinical outcomes with drug-eluting and bare-metal stents in patients with ST-segment elevation myocardial infarction: evidence from a comprehensive network meta-analysis. *J Am Coll Cardiol* 2013;62:496–504.
- Kim BK, Hong MK, Shin DH, *et al*; RESET Investigators. A new strategy for discontinuation of dual antiplatelet therapy: the RESET trial (Real Safety and Efficacy of 3-month dual antiplatelet Therapy following Endeavor zotarolimus-eluting stent implantation). *J Am Coll Cardiol* 2012;60:1340–8.
- Feres F, Costa RA, Abizaid A, *et al*. OPTIMIZE Trial Investigators. Three vs twelve months of dual antiplatelet therapy after zotarolimus-eluting stents: the OPTIMIZE randomized trial. *JAMA* 2013;310:2510–22.
- Gilard M, Barragan P, Noryani AA, *et al*. 6- versus 24-month dual antiplatelet therapy after implantation of drug-eluting stents in patients nonresistant to aspirin: the randomized, multicenter ITALIC trial. *J Am Coll Cardiol* 2015;65:777–86.
- Schulz-Schüpke S, Byrne RA, Ten Berg JM, *et al*; Intracoronary Stenting and Antithrombotic Regimen: Safety And Efficacy of 6 Months Dual Antiplatelet Therapy After Drug-Eluting Stenting (ISAR-SAFE) Trial Investigators. ISAR-SAFE: a randomized, double-blind, placebo-controlled trial of 6 vs. 12 months of clopidogrel therapy after drug-eluting stenting. *Eur Heart J* 2015;36:1252–63.
- Colombo A, Chieffo A, Frasieri A, *et al*. Second-generation drug-eluting stent implantation followed by 6- versus 12-month dual antiplatelet therapy: the SECURITY randomized clinical trial. *J Am Coll Cardiol* 2014;64:2086–97.
- Gwon HC, Hahn JY, Park KW, *et al*. Six-month versus 12-month dual antiplatelet therapy after implantation of drug-eluting stents: the efficacy of Xience/Promus Versus Cypher to Reduce Late Loss After Stenting (EXCELLENT) randomized, multicenter study. *Circulation* 2012;125:505–13.
- Valgimigli M, Campo G, Monti M, *et al*; Prolonging Dual Antiplatelet Treatment After Grading Stent-Induced Intimal Hyperplasia Study (PRODIGY) Investigators. Short-versus long-term duration of dual-antiplatelet therapy after coronary stenting: a randomized multicenter trial. *Circulation* 2012;125:2015–26.
- Fox KA, Carruthers KF, Dunbar DR, *et al*. Underestimated and under-recognized: the late consequences of acute coronary syndrome (GRACE UK-Belgian Study). *Eur Heart J* 2010;31:2755–64.
- Lee CW, Ahn JM, Park DW, *et al*. Optimal duration of dual antiplatelet therapy after drug-eluting stent implantation: a randomized, controlled trial. *Circulation* 2014;129:304–12.
- Collet JP, Silvain J, Barthélémy O, *et al*; ARCTIC investigators. Dual-antiplatelet treatment beyond 1 year after drug-eluting stent implantation (ARCTIC-Interruption): a randomised trial. *Lancet* 2014;384:1577–85.
- Mauri L, Kereiakes DJ, Yeh RW, *et al*; DAPT Study Investigators. Twelve or 30 months of dual antiplatelet therapy after drug-eluting stents. *N Engl J Med* 2014;371:2155–66.
- Navarese EP, Andreotti F, Schulze V, *et al*. Optimal duration of dual antiplatelet therapy after percutaneous coronary intervention with drug eluting stents: meta-analysis of randomised controlled trials. *BMJ* 2015;350:h1618–8.
- Bittl JA, Baber U, Bradley SM, *et al*. Duration of dual antiplatelet therapy: A systematic review for the 2016 ACC/AHA Guideline Focused Update on Duration of Dual Antiplatelet Therapy in Patients With Coronary Artery Disease. *J Am Coll Cardiol* 2016;68:1116–39.

- 30 Udell JA, Bonaca MP, Collet JP, *et al.* Long-term dual antiplatelet therapy for secondary prevention of cardiovascular events in the subgroup of patients with previous myocardial infarction: a collaborative meta-analysis of randomized trials. *Eur Heart J* 2016;37:390–9.
- 31 Farooq V, Serruys PW, Bourantas CV, *et al.* Quantification of incomplete revascularization and its association with five-year mortality in the synergy between percutaneous coronary intervention with taxus and cardiac surgery (SYNTAX) trial validation of the residual SYNTAX score. *Circulation* 2013;128:141–51.
- 32 G  n  reux P, Palmerini T, Caixeta A, *et al.* Quantification and impact of untreated coronary artery disease after percutaneous coronary intervention. *J Am Coll Cardiol* 2012;59:2165–74.
- 33 Yusuf S, Mehta SR, Chrolavicius S, *et al.* Fifth organization to assess strategies in acute ischemic syndromes investigators. Comparison of fondaparinux and enoxaparin in acute coronary syndromes. *N Engl J Med* 2006;354:1464–76.
- 34 Stone GW, Witzensbichler B, Guagliumi G, *et al.* HORIZONS-AMI Trial Investigators. Bivalirudin during primary PCI in acute myocardial infarction. *N Engl J Med* 2008;358:2218–30.
- 35 Rao SV, O'Grady K, Pieper KS, *et al.* A comparison of the clinical impact of bleeding measured by two different classifications among patients with acute coronary syndromes. *J Am Coll Cardiol* 2006;47:809–16.
- 36 Budaj A, Eikelboom JW, Mehta SR, *et al.* OASIS 5 Investigators. Improving clinical outcomes by reducing bleeding in patients with non-ST-elevation acute coronary syndromes. *Eur Heart J* 2009;30:655–61.
- 37 Karrow W, Vyas A, Giacomino B, *et al.* Radial versus femoral access for primary percutaneous interventions in ST-Segment elevation myocardial infarction patients. *JACC Cardiovasc Interv* 2013;6:814–23.
- 38 Valgimigli M, Gagnor A, Calabr   P, *et al.* MATRIX Investigators. Radial versus femoral access in patients with acute coronary syndromes undergoing invasive management: a randomised multicentre trial. *Lancet* 2015;385:2465–76.
- 39 Steg PG, Huber K, Andreotti F, *et al.* Bleeding in acute coronary syndromes and percutaneous coronary interventions: position paper by the working group on thrombosis of the european society of cardiology. *Eur Heart J* 2011;32:1854–64.
- 40 Mehta SR, Cannon CP, Fox KA, *et al.* Routine vs selective invasive strategies in patients with acute coronary syndromes: a collaborative meta-analysis of randomized trials. *JAMA* 2005;293:2908–17.
- 41 O'Donoghue M, Boden WE, Braunwald E, *et al.* Early invasive vs conservative treatment strategies in women and men with unstable angina and non-ST-segment elevation myocardial infarction: a meta-analysis. *JAMA* 2008;300:71–80.
- 42 Mart  nez-Sales V, Vila V, R  ganon E, *et al.* Elevated thrombotic activity after myocardial infarction: A 2-year follow-up study. *Haemostasis* 1998;28:301–6.
- 43 Trip MD, Cats VM, van Capelle FJ, *et al.* Platelet hyperreactivity and prognosis in survivors of myocardial infarction. *N Engl J Med* 1990;322:1549–54.
- 44 Hamid T, Zaman M. Switching of ticagrelor to Clopidogrel at 3 months in patients treated for acute care syndrome; Single centre experience. *Cardiovasc Pharm: Open Access* 2016;5.
- 45 Kerneis M, Silvain J, Abtan J, *et al.* Switching acute coronary syndrome patients from prasugrel to clopidogrel. *JACC Cardiovasc Interv* 2013;6:158–65.
- 46 Yeh RW, Secemsky EA, Kereiakes DJ, *et al.* DAPT Study Investigators. Development and validation of a prediction rule for benefit and harm of dual antiplatelet therapy beyond 1 year after percutaneous coronary intervention. *JAMA* 2016;315:1735–49.
- 47 Baber U, Mehran R, Giustino G, *et al.* Coronary Thrombosis and major Bleeding after PCI with Drug-Eluting Stents: Risk Scores from PARIS. *J Am Coll Cardiol* 2016;67:2224–34.
- 48 Breet NJ, van Werkum JW, Bouman HJ, *et al.* Comparison of platelet function tests in predicting clinical outcome in patients undergoing coronary stent implantation. *JAMA* 2010;303:754–62.
- 49 Gurbel PA, Erlinge D, Ohman EM, *et al.* TRILOGY ACS Platelet Function Substudy Investigators. Platelet function during extended prasugrel and clopidogrel therapy for patients with ACS treated without revascularization: the TRILOGY ACS platelet function substudy. *JAMA* 2012;308:1785–94.
- 50 Price MJ, Berger PB, Teirstein PS, *et al.* GRAVITAS Investigators. Standard- vs high-dose clopidogrel based on platelet function testing after percutaneous coronary intervention: the GRAVITAS randomized trial. *JAMA* 2011;305:1097–105.
- 51 The RISC Group. Risk of myocardial infarction and death during treatment with low dose aspirin and intravenous heparin in men with unstable coronary artery disease. the RISC group. *Lancet* 1990;336:827–30.
- 52 Lewis HD, Davis JW, Archibald DG, *et al.* Protective effects of aspirin against acute myocardial infarction and death in men with unstable angina. results of a veterans administration cooperative study. *N Engl J Med* 1983;309:396–403.
- 53 Cairns JA, Gent M, Singer J, *et al.* Aspirin, Sulfipyrazone, or both in unstable angina. results of a canadian multicenter trial. *N Engl J Med* 1985;313:1369–75.
- 54 White HD, French JK, Hamer AW, *et al.* Frequent reocclusion of patent infarct-related arteries between 4 weeks and 1 year: effects of antiplatelet therapy. *J Am Coll Cardiol* 1995;25:218–23.